

# *Tragopogon crocifolius*

## Habitus and growth type

Plant height [m]: **0.38**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **8.63**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Moisture indicator value: **2**

Reaction indicator value: **6**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.9**

Disturbance frequency (herb layer): **1.88**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.34**

Grazing pressure: **0.33**

Soil disturbance: **0.2**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**