

# *Tragopogon dasyrynchus*

## Habitus and growth type

Life span: **Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **5.5**

Moisture indicator value: **5**

Disturbance indicator values

Disturbance frequency: **1.83**

Disturbance frequency (herb layer): **2.11**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.54**

Mowing frequency: **0.85**

Grazing pressure: **0.38**

Soil disturbance: **0.24**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**