

# ***Tribulus terrestris***

## **Habitus and growth type**

Plant height [m]: **0.09**

Life span: **Annual**

Life form: **Therophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **43.48**

## **Flower**

Flowering period: **April-October**

## **Fruit, seed and dispersal**

Seed mass [mg]: **16.5**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

Environmental relationships

Substrate reaction relationship: **Alkaline**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**

Moisture indicator value: **3.5**

Reaction indicator value: **7.2**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.76**

Disturbance frequency (herb layer): **2.03**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.74**

Mowing frequency: **0.68**

Grazing pressure: **0.18**

Soil disturbance: **0.57**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*QD \(DIG\) \*Digitario sanguinalis-Eragrostietea minoris\*\*\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*V34 Trampled xeric grassland with annuals\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Synanthropic**