

# *Trifolium bocconeii*

## Habitus and growth type

Plant height [m]: **0.12**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **24.57**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.36**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **10**

Moisture indicator value: **3**

Reaction indicator value: **5.3**

Nutrient indicator value: **3.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.76**

Disturbance frequency (herb layer): **2.22**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.79**

Grazing pressure: **0.3**

Soil disturbance: **0.32**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*FH \(BUL\) \*Poetea bulbosae\*\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**