

Trifolium dalmaticum

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **7.4**

Moisture indicator value: **2.8**

Reaction indicator value: **6.5**

Nutrient indicator value: **2.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.34**

Disturbance frequency (herb layer): **2.08**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **0.51**

Grazing pressure: **0.36**

Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) *Sedo-Scleranthetea*](#), [EJ \(TRA\) *Stipo-Trachynietea distachyae*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**