

# *Trifolium pallescens*

## Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.62**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **2.8**

Moisture indicator value: **4.8**

Reaction indicator value: **4.7**

Nutrient indicator value: **2.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.36**

Disturbance frequency (herb layer): **1.63**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.03**

Grazing pressure: **0.22**

Soil disturbance: **0.15**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) \*Thlaspietea rotundifolii\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U71 Unvegetated or sparsely vegetated gravel bar in montane and alpine regions](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **6**

Continentality amplitude: **2**