

# *Trifolium pignanii*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **5.6**

Temperature indicator value: **6.7**

Moisture indicator value: **4.7**

Reaction indicator value: **6.3**

Nutrient indicator value: **4.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.01**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0**

Grazing pressure: **0.2**

Soil disturbance: **0.1**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T33 Mediterranean mountain Abies forest](#)

Broad habitat

Occurrence in broad habitats: **Forest**