

# *Tripidium ravennae*

## Habitus and growth type

Plant height [m]: **2.5**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **6.41**

## Flower

Flowering period: **June-October**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **9.3**

Reaction indicator value: **8**

Nutrient indicator value: **6.7**

Salinity indicator value: **2**

### Disturbance indicator values

Disturbance frequency: **0.87**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.22**

Grazing pressure: **0.21**

Soil disturbance: **0.19**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: **[FD \(NER\) Nerio-Tamaricetea](#), [MF \(JUN\) Juncetea maritimi](#)**

### EUNIS habitat

Diagnostic species of EUNIS habitats: **[N1J Mediterranean and Black Sea moist and wet dune slack](#)**

Constant species of EUNIS habitats: **[N1J Mediterranean and Black Sea moist and wet dune slack](#)**

Dominant species of EUNIS habitats: **[N1J Mediterranean and Black Sea moist and wet dune slack](#)**

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation, Scrub**