Tripolium pannonicum

Habitus and growth type

Plant height [m]: **0.45**
Life span: **Annual, Biennial or short-lived**
Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **17.96**

Flower

Flowering period: **July-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.91**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Reaction indicator value: **7.5**
Salinity indicator value: **7.2**

Disturbance indicator values

Disturbance frequency: **2.06**
Disturbance frequency (herb layer): **2.13**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.49**
Mowing frequency: **0.06**
Grazing pressure: **0.33**
Soil disturbance: **0.22**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **MA221 Atlantic saltmarsh driftline**, **MA223 Atlantic upper-mid saltmarsh and saline and brackish reed, rush and sedge bed**, **MA224 Atlantic mid-low saltmarsh**, **MA225 Atlantic pioneer saltmarsh**, **MA241 Black Sea littoral saltmarsh**, **MA252 Mediterranean upper-mid saltmarsh and saline and brackish reed, rush and sedge bed**, **R63 Temperate inland salt marsh**

Constant species of EUNIS habitats: **MA221 Atlantic saltmarsh driftline**, **MA223 Atlantic upper-mid saltmarsh and saline and brackish reed, rush and sedge bed**, **MA224 Atlantic mid-low saltmarsh**, **MA225 Atlantic pioneer saltmarsh**, **MA232 Baltic coastal meadow**, **MA241 Black Sea littoral saltmarsh**, **MA252 Mediterranean upper-mid saltmarsh and saline and brackish reed, rush and sedge bed**, **MA253 Mediterranean mid-low saltmarsh**, **N21 Atlantic, Baltic and Arctic coastal shingle beach**, **Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**, **R62 Continental inland salt steppe**, **R63 Temperate inland salt marsh**

Dominant species of EUNIS habitats: **MA224 Atlantic mid-low saltmarsh**, **R63 Temperate inland salt marsh**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Wetland, Saline vegetation, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **8**

Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz