

Typha domingensis

Habitus and growth type

Plant height [m]: **2.26**

Life span: **Perennial**

Life form: **Geophyte, Hydrophyte**

Leaf

Specific leaf area [mm²/mg]: **4.71**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.13**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.3**

Moisture indicator value: **10.3**

Reaction indicator value: **6.7**

Nutrient indicator value: **6**

Salinity indicator value: **1.5**

Disturbance indicator values

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **1.85**

Disturbance severity: **0.26**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.56**

Grazing pressure: **0.12**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[OD \(PHR\) Phragmito-Magnocaricetea](#)**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Sparsely vegetated (incl. rock and scree)**