Urtica pilulifera

Habitus and growth type

Plant height [m]: **0.58**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **January-June**

Fruit, seed and dispersal

Seed mass [mg]: **2.2**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**
Moisture indicator value: **5.4**
Reaction indicator value: **8**
Nutrient indicator value: **7.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.84**
Disturbance frequency (herb layer): **1.87**
Disturbance severity: **0.81**
Disturbance severity (herb layer): **0.7**
Mowing frequency: **0.41**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.24**
Soil disturbance: **0.37**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea*, QF (ART) *Artemisietea vulgaris***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T25 Phoenix theophrasti vegetation**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz