

# *Utricularia bremii*

## Habitus and growth type

Plant height [m]: **0.32**

Life span: **Perennial**

Life form: **Hydrophyte**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **12**

Reaction indicator value: **3.6**

Nutrient indicator value: **2**

Salinity indicator value: **0**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **[OB \(LIT\) Littorelletea uniflorae](#), [PA \(SCH\) Scheuchzerio palustris-Caricetea fuscae](#)**

Broad habitat

Occurrence in broad habitats: **Aquatic, Mire**

## **Distribution**

Continentality: **5**

Continentality amplitude: **3**