

Vaccinium macrocarpon

Habitus and growth type

Plant height [m]: **0.6**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Leaf

Specific leaf area [mm²/mg]: **12.59**

Fruit, seed and dispersal

Seed mass [mg]: **0.79**

Dispersal mode: **Endozoochory, Anthropochory**

Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Disturbance indicator values

Disturbance frequency: **1.14**

Disturbance frequency (herb layer): **1.25**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.01**

Grazing pressure: **0.35**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [PB \(OXY\) Oxycocco-Sphagnetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N1A Atlantic and Baltic coastal dune scrub](#)

Broad habitat

Occurrence in broad habitats: **Mire**

Distribution

Continentality: **5**

Continentality amplitude: **3**