

# *Valerianella carinata*

## Habitus and growth type

Plant height [m]: **0.2**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **32.97**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.55**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**

Moisture indicator value: **4.1**

Reaction indicator value: **7.6**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.17**

Disturbance frequency (herb layer): **1.96**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.57**

Mowing frequency: **0.45**

Grazing pressure: **0.29**

Soil disturbance: **0.42**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) \*Sedo-Scleranthetea\*, QC \(CHE\) \*Chenopodietea\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

## **Distribution**

Continentality amplitude: **6**