

Ventenata dubia

Habitus and growth type

Plant height [m]: **0.36**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**

Moisture indicator value: **3.3**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.59**

Disturbance frequency (herb layer): **1.71**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.38**

Grazing pressure: **0.29**

Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) Sedo-Scleranthetea](#), [QA \(PAR\) Papaveretea rhoeadis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

Distribution

Continentality: **6**

Continentality amplitude: **6**