

# *Veratrum nigrum*

## Habitus and growth type

Plant height [m]: **1.01**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **6.09**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5.6**

Temperature indicator value: **6.2**

Moisture indicator value: **4.3**

Reaction indicator value: **7**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.11**

Disturbance frequency (herb layer): **1.37**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.19**

Mowing frequency: **0.07**

Grazing pressure: **0.2**

Soil disturbance: **0.11**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) \*Carpino-Fagetea sylvaticae\*](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

## Distribution

Continentality: **7**

Continentality amplitude: **5**