

# *Veronica urticifolia*

## Habitus and growth type

Plant height [m]: **0.41**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **3.6**

Temperature indicator value: **3.8**

Moisture indicator value: **5.5**

Reaction indicator value: **7**

Nutrient indicator value: **6.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.03**

Disturbance frequency (herb layer): **1.25**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0**  
Grazing pressure: **0.19**  
Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [BA \(PIC\) Vaccinio-Piceetea](#), [CA \(FAG\) Carpino-Fagetea sylvaticae](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T31 Temperate mountain Picea forest](#), [T32 Temperate mountain Abies forest](#)

Constant species of EUNIS habitats: [T31 Temperate mountain Picea forest](#), [T32 Temperate mountain Abies forest](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

## Distribution

Continentality: **5**  
Continentality amplitude: **3**