

Vicia articulata

Habitus and growth type

Plant height [m]: **0.38**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **25.51**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **41.58**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **7.7**

Moisture indicator value: **4**

Reaction indicator value: **6.5**

Nutrient indicator value: **6.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.69**

Disturbance frequency (herb layer): **1.85**

Disturbance severity: **0.6**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.64**

Grazing pressure: **0.29**

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) *Chenopodietea*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**