

# *Vicia dumetorum*

## Habitus and growth type

Plant height [m]: **1.13**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **35.36**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6**

Temperature indicator value: **5.4**

Moisture indicator value: **5.2**

Reaction indicator value: **7.2**

Nutrient indicator value: **4.4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.1**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.67**

Disturbance severity (herb layer): **0.17**

Mowing frequency: **0.06**

Grazing pressure: **0.19**

Soil disturbance: **0.11**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) \*Trifolio-Geranietea sanguinei\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Synanthropic**

## Distribution

Continentality: **5**

Continentality amplitude: **3**