

# *Vicia ervilia*

## Habitus and growth type

Plant height [m]: **0.36**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **18.93**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **42.78**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Archeophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **8.8**

Moisture indicator value: **3.7**

Reaction indicator value: **6.3**

Nutrient indicator value: **5.3**

Salinity indicator value: **0**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**