

# *Vicia sparsiflora*

## Habitus and growth type

Plant height [m]: **0.62**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-June**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6**

Temperature indicator value: **7**

Moisture indicator value: **4**

Reaction indicator value: **6.5**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.02**

Disturbance frequency (herb layer): **1.28**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0.02**

Grazing pressure: **0.2**

Soil disturbance: **0.1**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) \*Carpino-Fagetea sylvaticae\*](#),  
[CB \(PUB\) \*Quercetea pubescentis\*](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**