

Vicia sparsiflora

Habitus and growth type

Plant height [m]: **0.62**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **May-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6**

Temperature indicator value: **7**

Moisture indicator value: **4**

Reaction indicator value: **6.5**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.02**

Disturbance frequency (herb layer): **1.28**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0.02**

Grazing pressure: **0.2**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#),
[CB \(PUB\) *Quercetea pubescentis*](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**