

# *Viola rupestris*

## Habitus and growth type

Plant height [m]: **0.07**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Seed mass [mg]: **1.11**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **4.9**

Moisture indicator value: **3.2**

Reaction indicator value: **7.8**

Nutrient indicator value: **1.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.54**

Disturbance frequency (herb layer): **1.59**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.24**

Grazing pressure: **0.27**

Soil disturbance: **0.16**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [AB \(KOB\) \*Carici rupestris-Kobresietea bellardii\*](#), [BC \(BRA\) \*Brachypodio pinnati-Betuletea pendulae\*](#), [BD \(PYR\) \*Pyrolo-Pinetea sylvestris\*](#), [CO \(ERI\) \*Erico-Pinetea\*](#), [DA \(FES\) \*Festuco-Brometea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H \*Larix light taiga\*](#)

Constant species of EUNIS habitats: [T3H \*Larix light taiga\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Oromediterranean scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **6**

Continentality amplitude: **6**