

# *Vulpia bromoides*

## Habitus and growth type

Plant height [m]: **0.2**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **20.86**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.43**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate reaction relationship: **Acidic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.6**

Temperature indicator value: **7.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.61**

Disturbance frequency (herb layer): **2.12**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.44**

Mowing frequency: **0.65**

Grazing pressure: **0.34**

Soil disturbance: **0.34**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) \*Koelerio-Corynephoretea canescentis\*](#), [FI \(TUB\) \*Helianthemetea guttati\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#), [R24 Iberian summer pasture \(vallicar\)](#)

Constant species of EUNIS habitats: [R12 Cryptogam- and annual-dominated vegetation on siliceous rock outcrops](#), [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#), [R24 Iberian summer pasture \(vallicar\)](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**

Continentality amplitude: **4**