

Zostera noltii

Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Hydrophyte**

Leaf

Specific leaf area [mm²/mg]: **30.08**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **6**

Moisture indicator value: **11.7**

Reaction indicator value: **7.8**

Nutrient indicator value: **5.5**

Salinity indicator value: **8**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [MA \(ZOS\) Zosteretea](#)

Broad habitat

Occurrence in broad habitats: **Marine, Aquatic**

Distribution

Continentality amplitude: **9**