Populus x canadensis

Habitus and growth type

Plant height [m]:
Life span: **Perennial**
Life form: **Phanerophyte, Tree**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Dispersal mode: **Anemochory, Anthropochory**
Dispersal distance class: **5, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **neophyte**

Ecology

Disturbance indicator values

Disturbance frequency: **0.15**
Disturbance frequency (herb layer): **1.4**
Disturbance severity: **0.79**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.03**
Grazing pressure: **0.2**
Soil disturbance: **0.22**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **T1H Broadleaved deciduous plantation of non site-native trees**
Constant species of EUNIS habitats: **T1H Broadleaved deciduous plantation of non site-native trees**
Dominant species of EUNIS habitats: **T1H Broadleaved deciduous plantation of non site-native trees**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Wetland, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz