

Anthemis aetnensis

Habitus and growth type

Plant height [m]: **0.16**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.51**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **3**

Moisture indicator value: **2**

Reaction indicator value: **1**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.92**

Disturbance frequency (herb layer): **0.92**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.8**

Mowing frequency: **0.02**

Grazing pressure: **0.15**

Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**FN \(RUM\) Rumici-Astragaletea siculi**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**U62 Mediterranean, Macaronesian and temperate volcanic field**](#)

Constant species of EUNIS habitats: [**U62 Mediterranean, Macaronesian and temperate volcanic field**](#)

Dominant species of EUNIS habitats: [**U62 Mediterranean, Macaronesian and temperate volcanic field**](#)

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**