

Bupleurum commutatum

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.9**

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **7.8**

Moisture indicator value: **3.1**

Reaction indicator value: **6.5**

Nutrient indicator value: **3.9**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **2.23**

Disturbance frequency (herb layer): **2.23**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.49**

Grazing pressure: **0.37**

Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#)

Broad habitat

Occurrence in broad habitats: **Scrub**