

Astragalus frigidus subsp. *frigidus*

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **23.57**

Fruit, seed and dispersal

Seed mass [mg]: **8.63**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **1.8**

Moisture indicator value: **4.5**

Reaction indicator value: **9**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.22**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.02**

Grazing pressure: **0.23**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) *Elyno-Seslerietea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U25 Boreal and Arctic base-rich scree and block field](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **8**

Continentality amplitude: **4**