# *Astragalus glycyphyllos* subsp. *glycyphyllos*

### Habitus and growth type

Plant height [m]: **0.68**
Life span: **Perennial**
Life form: **Hemicryptophyte**

### Leaf

Specific leaf area [$mm^2$/mg]: **25.66**

### Fruit, seed and dispersal

Seed mass [mg]: **4.93**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **6.5**
Temperature indicator value: **5.9**
Reaction indicator value: **7.1**
Nutrient indicator value: **3.4**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.18**
Disturbance frequency (herb layer): **1.39**
Disturbance severity: **0.64**
Disturbance severity (herb layer): **0.23**
Mowing frequency: **0.14**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Grazing pressure: **0.21**
Soil disturbance: **0.13**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R51 Thermophilous forest fringe of base-rich soils**
Constant species of EUNIS habitats: **R51 Thermophilous forest fringe of base-rich soils**

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

## Distribution

Continentality: **5**
Continentality amplitude: **4**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*