

# *Astragalus pelecinus* subsp. *pelecinus*

## Habitus and growth type

Plant height [m]: **0.14**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.79**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **10**

Moisture indicator value: **2.5**

Reaction indicator value: **7.5**

Nutrient indicator value: **3.5**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **1.75**

Disturbance frequency (herb layer): **2.14**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.48**

Grazing pressure: **0.4**

Soil disturbance: **0.31**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FH \(BUL\) \*Poetea bulbosae\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**