

# *Campanula sparsa* subsp. *sparsa*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Disturbance indicator values

Disturbance frequency: **0.35**

Disturbance frequency (herb layer): **1.45**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.22**

Grazing pressure: **0.24**

Soil disturbance: **0.17**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest](#)

Constant species of EUNIS habitats: [T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest](#)

Broad habitat

Occurrence in broad habitats: **Forest**