

# *Carlina macrocephala* subsp. *macrocephala*

## Habitus and growth type

Plant height [m]: **0.3**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **4**

Moisture indicator value: **3.5**

Reaction indicator value: **3.5**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.85**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.37**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.05**

Grazing pressure: **0.27**

Soil disturbance: **0.09**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FM \(PIL\) \*Saginetea piliferae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#)

Constant species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**