Cerastium fontanum subsp. *fontanum*

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Ecology

Disturbance indicator values

Disturbance frequency: **0.69**
Disturbance frequency (herb layer): **1.8**
Disturbance severity: **0.6**
Disturbance severity (herb layer): **0.49**
Mowing frequency: **0.48**
Grazing pressure: **0.19**
Soil disturbance: **0.19**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **U35 Boreal and Arctic base-rich inland cliff**
Constant species of EUNIS habitats: **U35 Boreal and Arctic base-rich inland cliff**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz