Cerastium fontanum subsp. *vulgare*

Habitus and growth type

Plant height [m]: **0.2**

Leaf

Specific leaf area [mm^2/mg]: **36.04**

Flower

Flowering period: **March-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.12**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Ecology

Disturbance indicator values

Disturbance frequency: **1.69**
Disturbance frequency (herb layer): **2.13**
Disturbance severity: **0.52**
Disturbance severity (herb layer): **0.51**
Mowing frequency: **0.99**
Grazing pressure: **0.33**
Soil disturbance: **0.36**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R21 Mesic permanent pasture of lowlands and mountains**, **R22 Low and medium altitude hay meadow**
Constant species of EUNIS habitats: **MA222 Atlantic upper saltmarsh**, **N18 Atlantic and Baltic coastal Empetrum heath**, **N1H Atlantic and Baltic moist and wet dune slack**, **R1S Heavy-metal grassland in Western and Central Europe**, **R21 Mesic permanent pasture of lowlands and mountains**, **R22 Low and medium altitude hay meadow**, **R23 Mountain hay meadow**, **R35 Moist or wet mesotrophic to eutrophic hay meadow**, **U71 Unvegetated or sparsely vegetated gravel bar in montane and alpine regions**, **V13 Arable land with unmixed crops grown by low-intensity agricultural methods**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Grassland (non-alpine, non-**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

saline), Sparsely vegetated (incl. rock and scree), Synanthropic

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz