

Halacsya sendtneri subsp. *devollensis*

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **6.1**

Moisture indicator value: **4.4**

Disturbance indicator values

Disturbance frequency: **1.69**

Disturbance frequency (herb layer): **1.69**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.42**

Grazing pressure: **0.31**

Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**