

# ***Hordeum marinum subsp. marinum***

## **Habitus and growth type**

Plant height [m]: **0.2**

Life span: **Annual**

Life form: **Therophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **45.54**

## **Fruit, seed and dispersal**

Seed mass [mg]: **4.15**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

### **Ellenberg-type indicator values**

Light indicator value: **8.9**

Reaction indicator value: **6.9**

Salinity indicator value: **4.7**

### **Disturbance indicator values**

Disturbance frequency: **1.48**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.17**

Grazing pressure: **0.3**

Soil disturbance: **0.27**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [JA \(SAG\) \*Saginetea maritimae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [MA251 Mediterranean upper saltmarsh](#), [R61 Mediterranean inland salt steppe](#)

Constant species of EUNIS habitats: [MA251 Mediterranean upper saltmarsh](#), [R61 Mediterranean inland salt steppe](#)

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Grassland (non-alpine, non-saline), Saline vegetation, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: **8**