

# *Paronychia chionaea* subsp. *chionaea*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **6.4**

Moisture indicator value: **2.8**

Reaction indicator value: **7.6**

Nutrient indicator value: **1.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.19**

Disturbance frequency (herb layer): **2.19**

Disturbance severity: **0.3**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0**

Grazing pressure: **0.3**

Soil disturbance: **0.13**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [\*\*KC \(ASP\) \*Asplenietea trichomanis\*\*\*](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**