

Salix cinerea subsp. *cinerea*

Habitus and growth type

Plant height [m]: **4.7**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **0.17**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Ecology

Disturbance indicator values

Disturbance frequency: **0.54**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.13**

Grazing pressure: **0.18**

Soil disturbance: **0.17**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S92 Salix fen scrub](#), [T15 Broadleaved swamp forest on non-acid peat](#), [T16 Broadleaved mire forest on acid peat](#)

Constant species of EUNIS habitats: [S92 Salix fen scrub](#), [T15 Broadleaved swamp forest on non-acid peat](#), [T16 Broadleaved mire forest on acid peat](#)

Dominant species of EUNIS habitats: [S92 Salix fen scrub](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**