

# *Salix cinerea* subsp. *oleifolia*

## Habitus and growth type

Plant height [m]: **4.7**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **12.3**

## Flower

Flowering period: **February-April**

## Fruit, seed and dispersal

Seed mass [mg]: **0.17**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Ecology

Disturbance indicator values

Disturbance frequency: **0.53**

Disturbance frequency (herb layer): **1.47**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.09**

Grazing pressure: **0.22**

Soil disturbance: **0.14**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FD \(NER\) Nerio-Tamaricetea](#), [HA \(POP\) Alno glutinosae-Populetea albae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S91 Temperate riparian scrub](#), [T14 Mediterranean and Macaronesian riparian forest](#)

Constant species of EUNIS habitats: [S91 Temperate riparian scrub](#), [T14 Mediterranean and Macaronesian riparian forest](#)

Dominant species of EUNIS habitats: [S91 Temperate riparian scrub](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**