Scrophularia ramosissima subsp. *macrorrhyncha*

Habitus and growth type

Plant height [m]: **0.3**
Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.3**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **9.5**
Moisture indicator value: **4**
Reaction indicator value: **6.5**
Nutrient indicator value: **4.5**
Salinity indicator value: **3.5**

Disturbance indicator values

Disturbance frequency: **1.75**
Disturbance frequency (herb layer): **1.88**
Disturbance severity: **0.39**
Disturbance severity (herb layer): **0.37**
Mowing frequency: **0.05**
Grazing pressure: **0.2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.41**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JD (AMM) *Ammophiletea arundinaceae***

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz