

# *Sedum ochroleucum* subsp. *mediterraneum*

## Habitus and growth type

Plant height [m]: **0.19**

Life span: **Perennial**

Life form: **Chamaephyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.04**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7.5**

Moisture indicator value: **2**

Reaction indicator value: **6**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.64**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.24**

Grazing pressure: **0.33**

Soil disturbance: **0.2**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) Koelerio-Corynephoretea canescentis](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#), [U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**