

# *Himantoglossum hircinum* var. *hircinum*

## Habitus and growth type

Plant height [m]: **0.39**

Life span: **Perennial**

Life form: **Geophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **20.02**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **7.2**

Moisture indicator value: **2.8**

Reaction indicator value: **8.6**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.94**

Disturbance frequency (herb layer): **1.73**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.66**

Grazing pressure: **0.31**

Soil disturbance: **0.15**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) \*Trifolio-Geranietea sanguinei\*](#), [DA \(FES\) \*Festuco-Brometea\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: **6**