

Lathyrus latifolius var. *latifolius*

Habitus and growth type

Plant height [m]: **1.45**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **18.7**

Fruit, seed and dispersal

Seed mass [mg]: **53.22**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**

Temperature indicator value: **7.5**

Moisture indicator value: **3.8**

Reaction indicator value: **8.3**

Nutrient indicator value: **3.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.37**

Disturbance frequency (herb layer): **1.52**

Disturbance severity: **0.6**

Disturbance severity (herb layer): **0.33**

Mowing frequency: **0.31**

Grazing pressure: **0.24**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) *Trifolio-Geranietea sanguinei*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S53 *Spartium junceum scrub*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

Distribution

Continentality: **4**

Continentality amplitude: **4**