

Trifolium echinatum var. *carmeli*

Habitus and growth type

Plant height [m]: **0.33**

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.94**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Moisture indicator value: **3.8**

Reaction indicator value: **2**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.41**

Disturbance frequency (herb layer): **1.92**

Disturbance severity: **0.77**

Disturbance severity (herb layer): **0.75**

Mowing frequency: **0.88**

Grazing pressure: **0.25**

Soil disturbance: **0.4**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**